


# WORLD'S WORST INSECT INVADERS: ANTS, BEES & WASPS



UPDATED: APRIL 2020

The Invasive Insects: Risks and Pathways project has found that the world's environmentally harmful invasive insect species are dominated by just one insect order – that of ants, bees and wasps (Hymenoptera)<sup>1</sup>. This order accounts for 16 of the 17 insect invaders known to be causing environmental harm in Australia. Why are ants, bees and wasps so successful and so harmful?

One answer is that they are habitual and versatile world travellers. Of the dozen main pathways by which invasive insects reach new countries, Hymenoptera use them all and frequently so<sup>1</sup>.

Another clue is that most invasive Hymenoptera are social, and the most harmful of them – typically ants – tend to live in extremely large societies, which can be more populous than the biggest human megapolises.

## DOMINANT INSECT INVADERS

Of the world's 24 insect orders, the Hymenoptera accounts for half the species in the assessment pool of the Invasive Insects: Risks and Pathways project (made up of species for which there is evidence of environmental harm somewhere in the world)<sup>1</sup>. The leading invaders are ants (all social), accounting for more than one in eight species assessed – three times as many as any other insect family. Bees, most of which are social, and parasitoid wasps, which are not, are also common invaders.

Of Australia's 17 insect invaders known to be causing environmental harm, 16 are Hymenoptera (9 ants, 3 wasps, 3 bees) and 1 is a beetle (Table 1)<sup>1</sup>. Three have been recorded in Australia for the first time just this century. As one indicator of the harm caused by social hymenopteran species, Australia is currently spending over \$60 million a year on eradication programs for five ant species – seeking



Western yellowjackets, invasive in Hawaii, are aggressive hunters.

Photo: TJ Gehling | Flickr CC BY-NC-ND 2.0



In their invasive range, European fire ants reach extremely high densities, displacing nearly all other ants.

Photo: Ryszard | Flickr | CC BY-NC 2

national eradication of red imported fire ants, electric ants and browsing ants, and partial eradication of yellow crazy ants (Wet Tropics) and Argentine ants (Norfolk Island).

The impacts of social hymenopteran species are complex, brought about mainly by predation, competition, and interaction with other exotic species<sup>1</sup>. Yellow crazy ants on Christmas Island demonstrate all these mechanisms. Capable of achieving extremely high population densities (more than 2000 a square metre), they aggressively displace

most other animals from their invasive range<sup>2,3</sup>. The ants benefit enormously by 'farming' an invasive scale insect from which they gain honeydew. They have killed tens of millions of red land crabs by spraying their eyes and mouthparts with formic acid. Because the crabs eat leaf litter, seeds and seedlings, their absence has drastically altered the structure and composition of invaded forests, and the forest canopy is suffering dieback due to the outbreaks of invasive scale insects protected by the crazy ants and from sooty mould caused by honeydew<sup>4</sup>. The loss of crabs also enables the spread of another invasive species, the giant African land snail<sup>5</sup>.

## WHAT WE MUST KEEP OUT OF AUSTRALIA

Of the 27 social hymenopteran species assessed so far in the Invasive Insects: Risks and Pathways project, 24 have been rated as 'of concern' or 'of substantial concern' (15 ants, 5 wasps, 4 bees)<sup>1</sup>. Of these, more than half (7 ants, 3 bees and 3 wasps) are already established in Australia. It is important to stop more introductions of these species – to

**Table 1: Invasive insect species for which there is evidence of environmental harm in Australia**

Order	Family	Species	Common name	Date of first detection or mention	First state or territory record
Hymenoptera	Apidae	<i>Apis mellifera</i>	European honey bee	1820	Queensland
Hymenoptera	Formicidae	<i>Solenopsis geminata</i>	Tropical fire ant	1863	?
Hymenoptera	Formicidae	<i>Paratrechina longicornis</i>	Black crazy ant	1886	Queensland
Hymenoptera	Formicidae	<i>Monomorium floricola</i>	Floral ant	1910	Queensland
Hymenoptera	Formicidae	<i>Monomorium destructor</i>	Singapore ant	1910	Queensland
Hymenoptera	Formicidae	<i>Pheidole megacephala</i>	African big-headed ant	1911	Queensland
Coleoptera	Scarabaeidae	<i>Heteronychus arator</i>	African black beetle	1920	South Australia
Hymenoptera	Formicidae	<i>Linepithema humile</i>	Argentine ant	1939	Victoria
Hymenoptera	Vespidae	<i>Vespula vulgaris</i>	Common wasp	1959	Victoria
Hymenoptera	Vespidae	<i>Vespula germanica</i>	European wasp	1959	Tasmania
Hymenoptera	Formicidae	<i>Anoplolepis gracilipes</i>	Yellow crazy ant	1975	Northern Territory
Hymenoptera	Vespidae	<i>Polistes chinensis</i>	Asian paper wasp	1979	New South Wales
Hymenoptera	Megachilinae	<i>Megachile rotundata</i>	Leafcutting bee	1987	New South Wales
Hymenoptera	Apidae	<i>Bombus terrestris</i>	Large earth bumblebee	1992	Tasmania
Hymenoptera	Formicidae	<i>Solenopsis invicta</i>	Red imported fire ant	2001	Queensland
Hymenoptera	Formicidae	<i>Wasmannia auropunctata</i>	Electric ant	2006	Queensland
Hymenoptera	Apidae	<i>Apis cerana</i>	Asian honey bee	2007	Queensland

prevent re-invasion of those being eradicated and new areas being invaded, and to stop new genetic material from boosting their invasiveness. This leaves at least 11 additional species that are likely to be a high priority to keep out of Australia (subject to Australia-specific risk assessment), including 8 ants, 2 wasps and 1 bee (see Table 2). There are likely to be other priority species once 28 additional Hymenoptera in the assessment pool have been assessed.

## WHY ANTS ARE SUCH SUCCESSFUL INVADERS

The Hymenoptera is one of, if not the largest of, insect orders, abundant everywhere except in polar regions and ecologically highly influential. A major reason for their success is the nutritious food they provide to their offspring<sup>6</sup>. Parental care in this order has been taken to a new level with the evolution of sociality, which has occurred several times, being the case for some bee and vespid wasp species and for all ants. Social structures enable dozens to millions of individuals to work cooperatively to exploit resources, build nests, care for young, and maintain strong defences.

**Table 2: The invasive social Hymenoptera not established in Australia that are of concern or substantial concern due to environmental impacts elsewhere in the world**

Family	Species	Common name
Apidae	<i>Apis mellifera scutellata</i>	East African lowland honey bee
Formicidae	<i>Myrmica rubra</i>	European fire ant
Formicidae	<i>Pachycondyla chinensis</i>	Asian needle ant
Formicidae	<i>Paratrechina fulva</i>	tawny crazy ant
Formicidae	<i>Solenopsis richteri</i>	black imported fire ant
Formicidae	<i>Technomyrmex albipes</i>	white-footed ant
Formicidae	<i>Lasius neglectus</i>	invasive garden ant
Formicidae	<i>Tapinoma melanocephalum</i>	ghost ant
Formicidae	<i>Solenopsis papuana</i>	Papuan thief ant
Vespidae	<i>Vespula pensylvanica</i>	western yellowjacket
Vespidae	<i>Vespa velutina</i>	Asian hornet

The most successful invasive Hymenoptera are the ants – which achieve an extreme form of sociality. A typical ant colony has one queen

attended by her sterile daughters, who aggressively defend their territory, particularly against members of their own species<sup>7</sup>. But many invasive ants



	Coleoptera	Dermaptera	Diptera	Hemiptera	Hymenoptera	Isoptera	Lepidoptera	Thysanoptera
<b>Biological control</b>	●				●		●	
Contaminant nursery material	●	●	●	●	●		●	●
<b>Food contaminant</b>	●		●	●	●		●	
Contaminant on plants	●		●	●	●		●	
Timber trade	●			●	●		●	
Transportation of habitat material	●	●	●		●	●		
Container/bulk	●	●	●		●		●	
Hitchhikers on ship/boat	●	●	●		●	●	●	
People and their luggage	●			●	●		●	
Vehicles	●			●	●	●	●	
Organic packing material	●	●			●		●	
Natural dispersal across borders	●		●	●	●		●	●

Figure 1: The use of introduction pathways by invasive insect orders. Only the most prevalent pathways and insect orders are shown. Circles and their sizes represent the relative contribution (%) of each insect order to the number of species using a particular pathway. Circle sizes (from smallest to largest) represent 1-10%, 11-20%, 21-30%, 31-40%, 41-50%, 51-60%.

form supercolonies, with many queens and many interconnected nests. The individuals from different nests behave amicably rather than aggressively towards each other and can move freely between nests. 'Thus, the cooperation that promotes the success of social insects generally appears to be exaggerated in many invasive ants'<sup>8</sup>.

The largest known supercolonies are those of the Argentine ant (*Linepithema humile*). In its native range, it usually forms colonies with small territories that are aggressively defended against other Argentine ants, and lives among many other native ant species<sup>8</sup>. Sometimes it forms supercolonies spanning hundreds of metres, although these endure for only

a few years, probably due to competition with other supercolonies and other ant species<sup>9</sup>. But their supercolonies in invaded areas can cover hundreds of kilometres, and some are more than a century old. One colony along the Mediterranean coast extends more than 6000 km, and one in Australia spans 2800 km<sup>9</sup>. These and several other supercolonies have their origins in just one ancestral supercolony from Argentina, and represent 'the most populous known animal society'<sup>10</sup>.

Normally, low genetic diversity is thought to be harmful but, for ants, by reducing their aggression towards their own kind, it frees up a lot of extra energy for colony growth, foraging, defence and competing

with other species<sup>7,8</sup>. Extremely high ant densities can be achieved, allowing domination over other species. Over 1 million Argentine ant queens and 4.4 cubic metres of workers and brood were captured from one 8-hectare orchard in the United States (cited in 10).

Ants also have flexible diets (as omnivores), easy-to-fulfil nesting requirements, and an association with humans that allows them to travel around the world in traded products and as stowaways on ships. The Invasive Insects Risks and Pathways project found they travel frequently on all 10 of the most prevalent unintentional pathways for insect introductions (see Figure 1).





Asian needle ant. Photo: © Matt Bertone

## IMPLICATIONS FOR BIOSECURITY

The prevalence and harmfulness of the invasive hymenopteran species, particularly ants, means they should be a top biosecurity priority in Australia. As tiny, tough and versatile travellers, they are also very challenging for biosecurity. We must strive to prevent new introductions, including of the species already here, eradicate those we can, and protect Australian biodiversity from their impacts.

### Recommendation 1

**Fully implement Australia's invasive ant biosecurity plan<sup>11</sup>.** This plan was adopted in mid 2019 by all national, state and territory governments but has no targeted funding for implementation. One particularly important action, essential for coordination and motivation, is to establish a 'permanent national body to coordinate national actions on invasive ants.' This is consistent with the seriousness of both existing and potential threats to the Australian environment, as well as human amenity and the economy.

### Recommendation 2

**Develop a biosecurity plan (or plans) for other social Hymenoptera.** With the potential for invasive bees and wasps to disrupt pollination communities, and the predatory efficiency of invasive wasps, they can have profound ecological impacts.

### Recommendation 3

**As a high biosecurity priority, reduce the risks of unintentional introduction pathways for the Hymenoptera.** Given their flexible travel habits – with ants in particular common on all 10 of the most prevalent unintentional pathways worldwide for invasive insects – the only feasible way to prevent new invasive hymenopteran species is to reduce the frequency by which they travel with traded products and as stowaways on ships and planes. This requires assessing the risks of each of these pathways, imposing import conditions to reduce the risks and improving inspection and diagnostic protocols.

### Recommendation 4

**As a high biosecurity priority, improve Australia's preparedness to respond to incursions of social Hymenoptera.** This requires developing effective surveillance programs and being ready to efficiently eradicate new incursions. Hymenopteran surveillance in Australia can be boosted by supporting citizen science programs.

## SOURCES

1. *Invasive Insects Risks and Pathways project unpublished data* (2019): Monash University and Invasive Species Council.
2. Abbott KL, Green PT (2005): Collapse of an ant-scale mutualism in a rainforest on Christmas Island. *Oikos*. 116(7): 1238-1246.
3. Abbott KL (2005): Supercolonies of the invasive yellow crazy ant, *Anoplolepis gracilipes*, on an oceanic island: Forager activity patterns,

density and biomass. *Insectes Soc.* 52: 266–273.

4. O'Dowd DJ, Green PT, Lake PS (2003): Invasional 'meltdown' on an oceanic island. *Ecology Letters*. 6: 812–817.
5. Green PT, O'Dowd DJ, Abbott KL, Jeffery M, Retallick K, Mac Nally R (2011): Invasional meltdown: invader–invader mutualism facilitates a secondary invasion. *Ecology*. 92: 1758–1768.
6. Quicke DLJ (2009): Chapter 127 - Hymenoptera: Ants, Bees, Wasps. In: Resh VH, Cardé RT, editors. *Encyclopedia of Insects (Second Edition)*. San Diego: Academic Press, pp 473–484.
7. Holway, Suárez ML, Case T (1998): Loss of intraspecific aggression in the success of a widespread invasive social insect. *Science*. 282: 949–952.
8. Tsutsui ND, Suarez AV (2003): The colony structure and population biology of invasive ants. *Conservation Biology*. 17: 48–58.
9. Vogel V, Pedersen JS, D'Ettoire P, Lehmann L, Keller L (2009): Dynamics and genetic structure of Argentine ant supercolonies in their native range. *Evolution*. 63: 1627–1639.
10. Van Wilgenburg E, Torres CW, Tsutsui ND (2010): The global expansion of a single ant supercolony. *Evolutionary Applications*. 3: 136–143.
11. Department of the Environment and Energy (2018): *Draft National Invasive Ant Biosecurity Plan 2018-2028*. Australian Government.

## ABOUT THIS PROJECT

The Invasive Insects: Risks and Pathways Project is a partnership between Monash University and the Invasive Species Council. To find out more visit [invasives.org.au/risks-and-pathways](https://invasives.org.au/risks-and-pathways).